



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Solution by G. I. HOPKINS, J. SCHEFFER, and G. B. M. ZERR.

Let AB be the sum of the two sides, CD the bisector, and F the given angle. Make $\angle HGK = \angle F$ and bisector $GL = CD$. Draw LM parallel to HG and NL parallel to GK . Extend GK making $GO = NG$. Make $OP = AB$, and on MP draw the semicircle MSP . Draw the perpendicular $MQ = ML$, also QR parallel to MP and ST perpendicular to MP . Through L draw TX , then TGX is the required triangle; for $TS^2 = MT \cdot PT = QM^2 = ML^2$. From similar triangles XNL and LMT , $XN:LM::NL:MT$. Since $MLNG$ is a rhombus $LM = NL$. $\therefore XN \cdot MT = ML^2$. $\therefore MT \cdot TP = MT \cdot XN$, whence $NX = TP$. $\therefore GX + GP = OP = AB$.

CALCULUS.

168. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, O.

The tangent of what Cartesian curve makes an x -intercept always m times as long as the corresponding y -intercept.

Solution by J. SCHEFFER.

Let the equation of the tangent be $y - y' = \frac{dy'}{dx}(x - x')$. Consequently the x -intercept $= x' - y' \frac{dx'}{dy'}$, and the y -intercept $= y' - x' \frac{dy'}{dx}$; therefore, omitting the accents, by the condition imposed upon the problem

$$-\frac{y}{p} + x = m(y - px); \quad \left(p = \frac{dy}{dx}\right),$$

whence $m x p^2 - (m y - x)p - y = 0$; or, arranged differently, $(px - y)(mp + 1) = 0$, whence $px - y = 0$ and $mp + 1 = 0$. From the former of these two equations we get $y = ax$; and from the second $my + x = b$, where a and b are arbitrary constants. Both equations represent straight lines, the first one of which passes through the origin.

MECHANICS.

159. Proposed by J. E. SANDERS, Hackney, Ohio.

Required the time for a tree, considered as a material line of uniform density, length $a = 100$ feet, to fall; the tree being inclined $\phi = 1^\circ$ from perpendicular.

Solution by the PROPOSER.

From Mechanics we find that

$$\frac{\sqrt{(2gh)t}}{l} = \int \frac{d\theta}{\sqrt{[1 - (2l/h)\sin^2 \frac{1}{2}\theta]}} \dots (1),$$